
# BIG BEND GAMBUSIA

# RECOVERY PLAN





1984

Albuquerque, New Mexico

# RECOVERY PLAN

FOR

# BIG BEND CAMBUSIA

Gambusia gaigei Hubbs, 1929

PREPARED FOR

THE U.S. FISH AND WILDLIFE SERVICE

BY

THE-RIO CRANDE FISHES RECOVERY TEAM

APPROVED: Letin X. Stanzan

Acting Regional Director, Region 2

ATE: 9/19

1. Point or condition when species will be considered for delisting:

The Big Bend gambusia may never be downliated or **celisted** because of its extremely limited distribution and **tenuous** habitat conditions. Downlisting the species to threatened **classification** could be considered if all recovery actions, a8 **proposed** In this Recovery Plan, are implemented.

2. What **must** be done to reach recovery:

Steps to reach recovery include maintaining and enhancing habitats where Big Bend <code>gambusia</code> presently exist and <code>establishing</code> populations in other <code>suitable</code> locations. To fnaure <code>againstloss</code> of <code>the</code> species a captive population must be maintained until survival of several population8 in the wild are assured.

3. Management needs to keep the species recovered:

To keep the species recovered it will be necessary to monitor population8 to insure they are viable and the habitats are being managed to maintain the species.

#### DISCLAIMER

This is the completed Big Bend Gambuaia Recovery Plan. It has been approved by the U.S. Fish and Wildlife Service. It does not necessarily represent official positions or approvals of cooperating agencies and It does not necessarily represent the **views** of all recovery team **members**/individuals, who played the key role in preparing the plan. This plan is subject to modification as dictated by new findings and changes in species status and completion of tasks described in the plan. Goals and objectives till be **obtained** and funds will be expended contingent upon appropriations, prioritea, and other **budgetary** constraints.

The Big Bend Gambuaia Recovery Plan dated **September**, 1984, **was** prepared by the U.S. Fish and Wildlife Service In cooperation with the Rio Grande Fishes **Recovery** Team.

The recovery plan is based upon the belief that State and Federal conservation agencies and knowledgeable, interested individuals should endeavor to preserve the Big Bend gambuaia and **its** habitat and to restore them, as much as possible, to their historic status. The objective of the plan is to make this belief a reality.

The recovery team members have **used** the beat information available to them and their collective knowledge and experience **in** producing **this** recovery plan. It is hoped the plan will be utilized by all agencies, **institutions**, and Individuals concerned with the Big Bend **gambusia** and the Rio Grande Village ecosystem to coordinate conservation activities. Periodically, and as the plan is implemented, **revisions** will be necessary. Revisions will be the responsibility of the recovery team and implementation is the task of the managing agencies, especially the **National** Park Service, Texas Parks and Wildlife **Department** and the U.S. Fish and Wildlife Service.

#### **ACKNOWLEDGMENTS**

The Big Bend Gambuala Recovery Plan, dated September 17, 1984, was prepared by the U.S. Fish and Wildlife Service in cooperation with the Rio Grande Fishes Recovery Team composed of the following individuals:

Dr. Clark Hubbs, Department of Zoology, University of Texas, Austin, TX Dr. Salvador Contreraa-Balderaa, Universidad Autonoma de Nuevo Leon, Monterrey, Mexico

Dr. Anthony Echelle, Biological Sciences, Oklahoma State University, Stillwater, OK

Mr. Buddy Jensen, Dexter National Fish Hatchery, Dexter, NM

Mr. Michael D. Hatch, New Mexico Department of Game and Fish, Santa Fe, NM

Mr. Floyd Potter, Texas Parks and Wildlife Department, Austin, TX

Other Individuals instrumental in plan formulations Include:

Mr. Gerard A. Hoddenbach, National Park Service, Santa Fe, NM

Mr. Gilbert Lusk, Park Superintendent, Big Bend National Park, TX

Mr. Mike Fleming, Resource Management Specialist, Big Bend National Park, TX

Mr. Bill Werrel, Water Resources Coordinator, National Park Service, Fort Collins, **CO** 

Literature citation should read as follows:.

U.S. Fish and Wildlife Service. 1984. Big Bend Gambusia Recovery Plan, U.S. Fish and Wildlife Service, Albuquerque, New Mexico.

Additional copies may be obtained from:

Fish and Wildlife Reference Service 1776 E. Jefferson Street, 4th Floor Rockville, Maryland 20852 Telephone: (301) 468-1737 Ext. 236 or 290 Toll Free - 1-800-582-3421

U.S. Fish and Wildlife Service Office of Endangered Species P.O. Box 1306 Albuquerque, New Mexico 87103 Telephone: (505) 766-3972

# TABLE OF CONTENTS

Summary*	i
Disclaimer	ii
Acknowledgments	iii
PART I - INTRODUCTION	
Introduction	1 1 2 4 5
Competition	5 6 7 8
Habitat Modification	9
Conservation Efforts	10
PART II <b>-</b> RECOVERY	
Objective	14
Stepdown Outline	14
Narrative	16
Literature Cited	30
PART III - IMPLEMENTATION SCHEDULE	32
PART IV - APPENDIX	
Replies to Comments	36
Monitoring Procedures	43

#### PART I

#### BIG BEND GAMBUSIA RECOVERY PLAN

#### INTRODUCTION

The Big Bend gambuaia (Gambuaia gaige1) la a livebearing fish, up to 30 mm long, restricted to one spring outflow near Rio Grande Village in Big Bend National Park, Brewster County, Texas.

Threats to the species include a very limited total environment, a declining spring flow that maintains the environment, and competition with <u>Gambuaia</u> affinia.

The species is a relatively plain, yellowish poeciliid whose faint lateral stripe is the most pronounced dark mark on the body. There la also a bar beneath the eye and a faint, dark chin bar. The male's anal fin is rolled into a tube-like gonopodlum for transferring sperm into the female, has a pronounced elbow, with only one or two segments. Hubbs and Springer (1957) give a &tailed description of the Big Bend gambuaia.


#### TAXONOMLG STATUS

The Big Bend gambusia was described by Hubbs (1929). Rosen and Bailey (1966) considered <u>Gambusia gaigei</u> a senior synonym of <u>Gambusia hurtadoi</u> and <u>Gambusia alvarezi</u>, but this opinion was not accepted by **Rivas**(1963), Peden (1973), Alvarez (1970), or Miller (1978); G. <u>gaigei</u> remains a distinct species.

Moat authors place the Big Bend gambuaia In the <u>Gambuaia nobilia</u> species group (Hubbs and Springer 1957, Minckley 1962, Peden 1973); however, Rivas (1963) divided the <u>G. nobilia</u> group Into the <u>G. nobilia</u> and <u>G. senilis</u> groups and placed G. <u>gaigei</u> in the latter.

#### DISTRIBUTION AND DESCRIPTION OF THE HABITAT

The Big Bend gambuaia is known only from spring habitats of Texas in the vicinity of Boquillaa Crossing and Rio Grande Village in Big Bend National The precise native occurrence is somewhat in question; the details are addressed in Hubbs, et al. (1977a). That study concluded that at least two populations originally existed--one at Boquillaa Spring approximately 200 meters north of Boquillas Crossing and a second at Spring 4 just east of the present Rio Grande Village Campground (Figure 1). Other names for Boquilias Spring are Boquillas Crossing and Beaver Marsh Springs, and other names for Spring 4 are Campground Springs (Brune 1981), Weden Spring, Graham Ranch Warm Springs, Rio Spring, and Pumphouse Big Bend gambusia possibly once existed in other springs in the Spring. vicinity of Rio Grande Village (notably Spring 1.) The population once inhabiting Boquillas Spring is extinct and the population in Spring 4 has been extirpated. The Big Bend gambuaia now consists or descendants of the Spring 4 population being maintained in a nearly natural park refugium (Fig. 2) and at Dexter National Fish Hatchery, New Mexico.






Figure 2. Rio Grande Village Campground showing location of Big Bend gambusia habitat.

The current stocks are all derived from three individuals taken from a declining Spring 4 population in 1956.

All documented natural occurrences of Big Bend gambusia are from spring outflows. The two springs of natural occurrence differ substantially in outflow temperature. Spring 4 has an outlet temperature of 32-35.5°C (Moore 1980) and Boquillas Spring had a temperature of about 23°C on March 6, 1982. However, the fish inhabiting Spring 4 did not live in the springhead itself (Hubbs 1959), but downstream where the temperature was influenced by ambient air temperatures. Recorded water temperatures (summers of 1954-56) from the Spring 4 outflow ditch, densely inhabited by Big Bend gambusia, were 28-32.5°C with air temperatures in the mid-20s. The temperatures in the large "overflow pool" (Figure 2), sparsely inhabited by Big Bend gambusia, were cooler and near ambient. Recorded temperatures from the refugium also fluctuate widely dependent upon air temperature and location.

The habitats originally occupied by the Big Bend gambusia were recorded as springfed marshes (Hubbs 1929). Dense aquatic vegetation (submerged and emergent) presumably occurred in clear, shallow water. Alt hough the present refugium has open water with depths in excess of one meter, the Big Bend gambusia is most abundant among the vegetation (primarily Typha and Charn) near the shore.

In summary, the natural habitat occupied by <u>G. gaigei</u> was clear, shallow water fed by warm springs. Although the Big Bend gambusia will exist in other habitats, the rapid replacement by mosquitof ish <u>(Gambusia affinis)</u> (see Threats) in eurythermal water suggests that the Big Bend gambusia

nobilis species group in U.S. waters also are abundant in springs, especially in those area6 inhabited by mosquitofish downstream.

#### ASSOCIATED SPECIES

Big Bend gambusia have been found associated with at least five other fish species within their native range. These included, at Spring 4, two minnows (Notropis braytoni and N.lutrensis), another poeciliid (Gambusia affinis), an ictalurid (presumably Ictalurus punctatus) and a centrarchid (Lepomis The channel catfish and green sunfish (potential predators) were reported to have been purposefully stocked into the overflow pool of Spring 4; perhaps the other specie6 accidentally were placed into the pool along with these game specie6 (Hubbs and Springer 1957, Hubbs and These predators were removed and their impacts reportedly Broderick 1963). were minimal (Hubbs and Echelle 1972). The two potentially competitive minnows also were eliminated, but the mosquitofish survived and apparently Ls a serious competitive and predatory threat to Big Bend gambusia. A second introduction of green sunfish occurred in 1968, apparently as an accident by campers who wished to "help the conservation program" (Hubbs and Echelle 1972). These also were eliminated with few noticeable effects on the Big Bend gambusia. At the present time, no fishes are known to coexist with Big Bend gambusia.

At least 18 other fish species inhabit the Rio Grande adjacent to the habitat6 of the Big Bend gambusia (Hubbs et al. 1977b) but are not syntopic with the Big Bend gambusia. They could move into the campground area in the event of a large-magnitude flood.

Other species found with Big Bend gambusia 'include aquatic invertebrate6 that may be prey items for G. gaigei, crayfish, a variety of potential avian predators, and various plant6 including bulrushes, Cattails, saltcedars, and cottonwoods, which provide cover and/or shade.

#### REPRODUCTION

The Big Bend gambusia is viviparous (bear6 living young). Hubbs and Broderick (1963) report two broods totalling 40 young from two small females. The species reproduces in laboratory tanks and at the Dexter National Fish Hatchery in ponds fed by well water. Otherwise, virtually nothing is known about reproduction in the species. However, it is probable that, once inseudnated, females can store sperm for long periods, e.g., several months, and that interbrood intervals in individual females is on the order of 1-2 months. These are general features of reproduction in Gambusia species. Factors limiting reproduction are unknown, but photoperiod, temperature, and food availability all affect fecundity and interbrood intervals in poeciliids.

#### COMPETITION

The evidence that competition with G. affinis is deleterious to

the survival of Big Bend gambusia population6 is primarily circum-Other factors such as predation on young may be involved stantial. (Meffe 1983). Also, Gambusia gaigei does not flourish in the presence of Gambusia affinis (Hubbs et al., 1977a). The recorded evidence of this includes: (1) the replacement of Gambusia gaigei by Gambueia affinis in the Spring 4 outflow (1954-56); (2) replacement of Gambusia gaigei by Gambusia affinis in the first refugium (Fig 2; 1960-1961); (3) the failure of reintroduced Gambusia gaigei to establish a population in Boquillas Spring where Gambueia affinis now thrive; (4) the rapid replacement of Gambusia gaigei by Gambusia affinis in artificial pools near Brownsville. This last example included disappearance of mature Gambusia gaigei within 3 months of the first appearance of Gambusia affinis suggesting that c&petition or predation on adults as well as the young. In general, the rates of replacement were most rapid in eurythermal habitats (fluctuating temperatures) and slowest in warm, springrun habitats (constant temperatures).

#### DEVELOPMENT

Visitors to Big Bend National Park continue to increase with coincidental impacts on the park's natural resources. In the future, satisfaction of additional visitor needs will require careful planning to avoid overburdening the Rio Grande Village area with new demand.6 for water and space. Current visitor use of the Rio Grande Village area may be approaching the maximum commensurate with management for survival

of the Big Bend gambusia, with the main threat being increased pressure on the park to provide additional facilities. Water supply for Rio Grande Village presently is pumped from Spring 4, utilizing most of the natural flow from this spring.

#### RECHARGE ZONE

Spring 1 and Spring 4 are Important for sustaining <u>Gambus1a gaigei</u> in the Rio Grande Village area of Big Bend National Park. These springs (about 270 meters apart) have been developed, altering their natural flows. Chemical water analyses show, them to have been almost identical during the past 24 years and strongly suggest that both springs have a common source. Spring6 1 and 4 are intermediate thermal springs, part of acluster of six known thermal springs concentrated in this area of the Big Bend of the Rio Grande. Temperatures of the spring cluster are in the 30–50°C range with the warmest temperatures in Hot Springs and temperatures in Springs 1 and 4 near 34°C. The points of issue of Springs 1 and 4 are located along the contact of the upthrow side of the Santa Elena Limestone of Cretaceous Age (Moore 1980).

Thermal water issuing from both springs is of ancient origin with the principal. recharge zone located in higher elevations of the Sierra del Carmen (Moore 5.980). The fault system, formed in Mesozoic times during development of the Sierra del Carmen, serves both as a means for collecting and recharging the groundwater system as well

as an avenue by which groundwaters are discharged In a spring line. In addition to the concentrated flows of Springs 1 and 4, diffused seepage occurs in the vicinity.

No land-use changes or significant water developments are anticipated over the recharge zone; however, monitoring of Spring 1 and Spring 4 discharges should continue so that potential changes in water quality or quantity are detected as early as possible.

## RUNOFF

Surface runoff and Rio Grande flooding are two continual threats to the existence of Cambusia gaigei. Periods of unusually high precipitation could result in surface runoff causing significant silt deposition and/or erosion in both the refugium and/or springruns. Flooding or broad surface runoff could provide an avenue for invasion of Gambusia gaigei habitats by Gambusia affinis (or other undesirable species) and renew the threat of biological contamination. The greater of the two threats is flooding by the Rio Grande. Flow in the Rio Grande near Big Bend is regulated primarily by water released from Presa Luis L. Leon into the Rio Conchos in Mexico. Cooperative efforts should be maintained with the Republic of Mexico to prevent water releases Large enough to flood Rio Grande Village. However, high rainfall and seasonal runoff in the Rio Grande are not predictable nor fully controllable and could result in flooding of Gambusia gaigei habitat.

#### **HABITAT MODIFICATIONS**

The history of spring visitation by scientists interested in observing and collecting Big Rend gambusia is sketchy and has led to considerable discussion about which spring is the type locality and which springs were once occupied by the species. This is covered in detail by Hubbs et al. (1977a).

Available evidence is that the type locality (Boquillas Spring) dried during the 1950s. Although flows have now resumed, the waters are occupied by dense populations of <u>Gambusia affinls</u>. Thus, inis present condition, the type locality is no longer a favorable habitat for Big Band gambusia. The other actual or potential habitats for Big Bend gambusia are the outflows of Spring 1 and Spring 4.

Spring 4 originally flowed almost due west about 20 m. ters and then southwest for about SO meters to a conjunction with runoff draining the northeast side of the Rio Grande Village valley (Fig. 2). The joined streams flowed about 200 meters south to the Rio Grande. In 1951, the flow was diverted by a dike so that it flowed directly south into a Large overflow pool. This pool is now a choked, cattil slough; excess water flows south to join the old channel just prior to entry into the Rio Grande. Presently, some water flows through both courses; the original flow passes through a wooded grove until it crosses the nature trail. Beyond that point, it enters another cattail slough associated with a beaver pond. The diverted part of the streamisfilled with cattails

until It crosses the nature trail (the trail occupies &he crest of the dam forming the overflow pool). Most of the water from Spring 4 is used as a water supply for Rio Grande Village.

Little is known concerning the early **history** of Spring 1. Early maps suggest an outflow to the east toward Berkeley House. Other changes in the area are **discussed** under conservation efforts.

#### CONSERVATION EFFORTS

Over the years, there have been projects designed to rodify spring outflows through ditches and ponds to enhance the, appearance of the "oasis image" of the Rio Grande Village area. Since the 1950s, these projects have been cognizant of Big Bend gambusia requirements.


Chief conservation efforts have been to ensure the survival of the Spring 4 population. The various problems are treated In Huibs and Broderick (1963), Wauer (1973), Hubbs et al. (1977a), Hubbs and Williams (1979). The initial difficulties occurred in 1954 through 1955 when Gambusia affinis was observed to be replacing Big Bend gambusia in Spring 4. The October 9, 1956, renovation involved an unsuccessful of fort to eradicate Gambusia affinis from Spring 4. On that date, live B.g Bend gambusia were removed from the Spring 4 outflow. Subsets of the Live sample were

introduced Into potentially suitable Park Service waters. All of these fish died; the only remaining living individuals were three fish maintained In Austin and returned to the park the following spring (1957). These fish were stocked into the first refuglum pond built during the 1956-57 winter. Subsequently, Gambusia affinis were noted to be abundant in the refugium pond. On April 16, 1960, 15 Big Bend gambusia were removed from the initial refugium and taken to Austin. During the summer, the current refugium near Spring 1 was constructed and filled with Spring 4 water. Twenty fish were placed in this refuglum August 7, 1960.

During 1968, green sunfish **were** unexpectedly observed in the new refugium pond and these predators were eliminated.

A massive mortality of 3ig Bend gambusia in the refugium was noted in December 1975. The observations are in accord dth cold shock le thali ty (Hubbs and Williams 1979). In August 1976, the few remaining fish in the refugium were supplemented by descendents of the 1960 stocks that had been maintained in Austin.

In late 1976, a constant flow-through system based on pumped water from Spring 1 was established in the refugium. This system was designed to mitigate the effects of thermal extremes such as those that presumably caused the December 1975 mortality. Overflow water drained to the south and into a saltcedar/cattail thicket. The overflow was er was concentrated in an overflow ditch (Fig. 3). In January 1978, Big B. nd gambusia from



the refuglum were placed in the overflow drainage ditch and they now flourish In the refuglum and in its drainage ditch.

A stock of Big Bend gambusia was first taken to Dexter National Fish Hatchery in 1974. Low winter temperatures eliminated <code>several</code> stockings at Dexter, but present procedures call for <code>maintaining some</code> fish in warmer hatch-house waters throughout the tinter. This change has allowed the present population to survive since 1981, and guarantees replacement of Big Bend gambusia in the event of a massive <code>mortalicy</code> of the species in Big Bend National Park.

Future management goals for recovery of the species include the supplement of spring flows from wells, extending the outflow from Spring 1 to form a slow, shaded creek, redirecting the flow from Spring 4 to form a stream which will preximate predevelopment conditions, eradication of G. affinis from the campground area, and the establishment of Big Bend gambusia in other suitable locations.

#### RECOVERY

# Objective

The ultimate goal of this recwery plan is to secure survival of the Big Rend gambusia in a natural setting. This goal **should** result from implementation of the recovery plan proposed below. because of the extremely limited distribution and tenuous habitat conditions, Big Rend gambusia may never be downlisted or at least delisted.

# Step-Down Outline

Primary goal: Assure the survival of the Big Bend gambusia (Gambusia gaigei) through improvement of its status.

- 1.0 Maintain and enhance the existing Big Bend gambusia population and Its habitat.
  - 1.1 Identify population needs.
    - 1.11 Competition with Gambusia affinis
    - 1.12 Prey species biology
    - 1.13 Reproductive variables
    - 1.14 Predation
    - 1.15 Surviorship
    - 1.16 Diseases and parasites
    - 1.17 Habitat requirements

- 1.2 Monitor existing population and habitat.
- 1.3 Evaluate, maintain, and enhance habitat.
  - 1.31 Supplement spring flow
  - 1.32 Manage Spring 1
  - 1.33 Manage Spring 4
  - 1.34 Irrigate campground from a well
  - 1.35 Eradicate Gambusla affinis from campground area
  - 1.36 Minimize campground impact
- 2.0 Establish Big Bend gambusia in suitable locations.
  - 2.1 Determine suitable locations . .
    - 2.2 Stock Big Bend gambusia in Spring 4
    - 2.3 Use Big Bend gambusia for mosquito control in the Big Bend area
    - 2.4 Monitor released populations
    - 2.5 Manage Introduced populations
- 3.0 Maintain a captive population of Big Bend gambus: a.
- 4.0 Produce information for public consumption.
  - 4.1 Information phamplet
  - 4.2 Interprecive programs
  - 4.3 News releases
- 5.0 Enforce Federal and State laws.
  - 5.1 Status
  - 5.2 Habitat integrity

#### Narrative

Recovery Goal: Assure the survival of the Big Bend gambusia (Gambusia gaigei) through improvement of its status.

1.0 Maintain and enhance the existing Big Bend gambusia population and its habitat.

The only known natural population of Big Bend gambusia inhabits a refugium in Big Bend National Park. The recovery team recommends that the first priority for recovery of the species be enhancement of suitable habitat. Unless the environment is appropriately modified to assure continuity, survival of the species remains under continued threat. In addition, the exceedingly small geographic rang is of great concern because one seemingly minor environmental change could exterminate the species from the wild.

#### 1.1 Identify population needs

It is essential to understand the biology and ecology of Big Bend gambusia. In the event the status of the species declines, causative factors can be identified and remedied as quickly as possible.

- 1.11 Assess potential for competition with Gambusia affinis.

  Gambusia affinisis a known and continuing potential competitor with G. gaigei. It is essential to understand competitive interactions between Big Eend gambusia and G. affinis because of the serious and persistent threat the latter species poses to the former. Studies should be conducted to better understand the competitive interactions for food and space between these two species.

  Contrast the biological requirements of each species whereever appropriate.
- 1.12 Determine food preferences and prey biology.

Inventory the prey species present in the refugium and in potential restoration sites. In addition, conduct studies to determine food preferences of Big Bend gambusia and seasonal food availability.

1.13 Determine reproductive variables of Big Bend gambusia.

Additional knowledge of the reproductive biology (reproductive season, fecundity, interbrood Interval) of the species is not urgent to this recovery plan but studies should be pursued wherever possible to develop a reproductive data base useful to enhance perpetuation of the species.

1.14 Determine effects of predation.

Lepomis cyanellus has been linked to a decline of Big Bend gambusia during the 1960s. Letermine effects of predation by potential predators (birds, sunfish, etc.). This study should involve selectivity on Big Bend gambueia by predators.

1.15 Determine survivorship by age groups.

Little is known about survivorship curves for Big Bend gambueia. Mortality rates for each life history stage should be determined and that information incorporated into a plan for reducing mortality.

1.16 Determine diseases and parasites.

No data are available on the diseases and parasites .

affecting Big Bend gambusia. As the species occupies

limited space, an epidemic could seriously impact chances of survival. Advance knowledge of the diseases and parasites of Big Bend gambusia could be of assistance in containing an epidemic.

#### 1.17 Determine habitat requirements.

Valuable Insight for protection and enhancement of Big
Bend gambusia would be **gained** from a study of seasonally
varying habitat associations including physical, chemical,
and biotic **elements** of present habitat and potential
restoration sites.

Develop an adequate and dependable water source to satisfy the basic habitat requirement of Big Bend gambusia. Diminished spring cutflows have resulted in loss of some of the original Big bend gambusia habitat. The human demand for water in the park is likely to exacerbate the problem. Spring flows in the refugium and potential restoration sites must be enhanced.

# 1.2 Monitor Big Bend gambusia and its habitat.

Conduct long term population and monitoring studies of Big Bend gambusia focusing interests on habitat conditions, population numbers, condition, and age structure of fich. Should any of these or other factors suggest decline in the population or degradation of habitat, causative factors must be identified and corrected. Every proposed activity which might adversely affect the Big Bend gambusia or its habitat must be critically reviewed.

#### 1.3 Evaluate, maintain, and enhance habitat.

Big Bend gsmbusia survivorship **18** dependent upon availability of a warm spring habitat. The species survives well in other habitats only in the absence of <u>G. affinls</u>; but the adverse effect of <u>G. affinis **18**</u> least important in warm springrun environments.

#### 1.31 Supplement spring flow.

(Springs 1 and 4) have adequate available outflow volumes for stable Big Bend gsmbusia populations. The ground water in the vicinity of the Rio Grande Village Campground emerges at several locations. Wells should be dug into the aquifer upstream from the outflows of Springs 1 and 4. Water from these wells should be available to supplement the flows from Springs 1 and/or 4 when either flow is diminished. Each spring (see 1.32 and 1.33 below) should have consistent flow to maximize the stenothermal warm environment most optimal for Gambusia gaige1 populations. Flow rate is less critical than maintenance of stenothermal conditions typical of springrun habitats.

#### 1.32 Manage Spring 1.

Spring 1 is used to provide water for the existing refugium. The outflow from the pool now drains into a saltcedar/cattail thicket. This outflow should be extended along the present asphalt road in the form of a slow, shaded creek. No water should be permitted to reach the Rio Grande. It should be dispersed on flatlands in the vicinity of the maintenance yard. The vegetation developing along the slow creek should be monitored carefully. Cattails and bullrushes should be removed to preclude overgrowth. Appropriate native shade trees should be encouraged.

#### 1.33 Manage Spring 4.

The outflow from Spring 4 contained the ancestors of the present stocks of Big Bend gembusia, and some individuals may still exist in the dense vegetation. Excess water not needed for the fish could be used for domestic use in the campground, but priority should be placed on habitat for the endangered species. All water for fish use should be directed into the original channel. Redirection of flow can be achieved by removal of a portion of the existing dike blodcing flow towards the old channel and placing dirt across the ditch carrying water to the overflow pool. Concentration of the water in the channel that flows across the nature trail will recreate a stream that approximates presumed predevelopment circumstances.

The flow in the **combined** stream should **follow** the natural stream course. The stream should have a consistent flow with minimal **shallow**, still water; however, the flow should not be directed through a straight channel. Factors **that** might cause ponding of the stream (beaver activities, cattail growth, etc.) should be prevented.

1.34 Irrigate the campground from a well.

The present practice of using water directly pumped from the Rio Grande to water vegetation in Rio Grande Village should be discontinued to avoid contamination with **G**. afflnis. A shallow well should be dug into the river gravel in a secure location at least **50** meters from the river bank. This **well** (or wells) should be the source of irrigation water for the campground.

1.35 Eradicate Gambusia affinis from campground area.

Any population of Gambusia affinis near waters inhabited by Gambusla gaigei is a potential source of an introduction; thus, all likely sources of contaudnation should be removed. Populations of G. affinis exist In most exposed open water in the vicinity of Rio Grande Village. Those populations should be eliminated. Many of the populations can be elindnated by drying of the pools. Use of irrigation

water (from the well(s) described In 1.34) should be monitored carefully so water does not accumulate in low spots. Such inadvertant pooling should be allowed to dry so that a crust forms across the bottom; however, subsequent to canplete drying, overflow water may be allowed to accumulate In these ponds. When G. affinis has been eradicated frw all poads in and adjacent to the campground, the only sources of potential contamination will be populations in the Rio Grande and in the outflow from Spring 4. It is not fess ible to eradicate G. affinis from the Rio Grande and past attempts in Spring 4 have been unsuccessful; however, any activities that enhance G. affinis abundance in either area should be avoided.

#### 1.36 Minimize campground Impact

Any activities at the Rio Granda Village Campground that adversely impact Big Bendgambusia must be avoided.

Campsites close to Splng 1 or 4 outflows till be reevaluated. Nearby campsites should be closed and new or replacement campsites be located a greater distance from the springruas or sources. Alternate camping localities elsewhere in the park should be considered to protect Gambusia gaigai habitat.

# 2.0 Establish Big Bend gambusia in suitable locations.

Only warm spring habitats within the Boquillae Spring - Rio Grande Village area (Figure 1) are suitable for introduction of Big Bend gambusia.

Purposes of this action are to: (1) increase the numbers of viable stocks within the historic range of the species; (2) provide transplantation experiments in a variety of situations and thereby gain increased understanding of the breadth of conditions tolerated by the species; (3) provide situations where <code>genetic/ecological</code> interactions with <code>G</code> affinis can be exandned in natural conditions; and (4) reduce the size of the resident G. affinis population, thereby reducing the threat to <code>G</code> gaigei.

In the long run, <u>G. affinis</u> will be a persistent **component** of the local fauna with which <u>G. gTigle1</u>emust contende q u e n c e s of interaction between the **two** species must be understood.

## 2.1 Determine suitable locations to stock Big Bend gambusia

Spring 4 is the only warm spring habitat currently suitable for <u>G. gaigei</u>, however, ponds in the campground may also be suitable.

# 2.42 Establish emergency response procedures

When routine monitoring reveals a degradation of the status of Big Bend gambusia or habitat, procedures for assuring survival of the species should be developed and ready to be put into affect.

# 2.5 Manage introduced populations

## 3.0 Maintain a Captive Population of Big Bend Gambusia

Big Bend gambusia should be maintained at Dexter National Fish Hatchery to provide a'resgrve gene pool in the event that the species experiences a severe population decline in its natural setting. Standard procedures at the hatchery should be followed. This facility has the capability to safely overwinter <u>G. gaigei</u> and to isolate the species from other gambueine fishes. The captive populations can be used to provide research specimens (both live and preserved) to agencies and institutions conducting research on G. <u>gaigei</u>. If needed, designated studies can also be carried cut at the holding facility. The stock at Dexter should be checked annually to verify the genetic integrity of the captive population.

#### 4.0 Information Base

Funds and manpower should be expended to provide public information on survival problems and recovery efforts relevant to Gambusia gaigei in Big Bend National Park.

# 4.1 Information pamphlet

Information relative to the **uniqueness,taxonomy,** biology, distribution, and habitat requirements of the Big Bend gambusia should be prepared and made available to enhance public awareness of endangered species in general, and this species in particular.

# 4.2 Interpretive Programs

Information derived from such pamphlet should be presented to the visiting public in park interpretive programs and in visitor center displays.

#### 4.3 News Releases

Noteworthy events regarding the Big Bend gambusia should be publicized through the dissemination of appropriate and timely news releases.

#### 5.0 Law Enforcement

The Big Bend gambusia is currently protected under Federal and Texas laws. It occurs only in Big Bend National Park where all natural resources are federally protected and where National Park Service law enforcement personnel are readily available. State and Federal enforcement agencies and divisions of the National Park Service

# 2.2 Stock Big Bend gambusia in Spring 4

Once actions described in 1.33 have been implemented, Big Bend gambusia should be transported from Spring 1 refugium to the upper portion of the Spring 4 run. Transplantation of at least 100 females and 100 males is recommended. Repeated efforts should be made if initial introduction(s) are unsuccessful.

Spring 4 should provide a valuable opportunity to examine resource utilization differences and genetic interaction between G. gaigei and G. affinis.

#### 2.3 Use Big Eend gambusia for mosquito control in Big Bend area.

At present the park maintains a large pool which receives overflow irrigation water from the campground area (Figure 1). This pool represents a large reservoir of <u>G. affinis</u> which is a threat to the well being of Big Bend gambusia. This pool should be stocked with <u>Gambusia gaigei</u> after Implementation of 1.35.

2.4 Monitor relocated populations of Big Bend gambusia.

To assess the **success** of transplantations and other management actions, all known populations at Big Bend should be sampled at least twice **annually.** Such monitoring should be **coupled** with assessments of habitat quality.

2.41 Establish monitoring procedures and schedules

Sample all sites twice a year, in February and August.

Extreme care should be taken to guard against inadvertent transport of eggs/embryos of **G. affinis** and other fishes.

The procedures for sampling is appended to this plan.

- 2.411 Preset.;: refugium should sample, preserve, and catalogue at least 15 specimens, unless such sample represents more than 1 percent of the standing populations numbers.
- 2.412 Similar sampling in the overflow stream from

  Spring 4 done by sources selected by the National

  Park Servive in consultation with the Fish and

  Wildlife Service.

should be provided information relevant to Big Bend gambusia identification, legal status, distribution and maintenance. Habitat integrity should be emphasized to avert actions (intentional or unintentional) or projects that would be deleterious.

#### 5.1 Status

Enforcement agencies (Federal and State) will be kept advised of the legal status of the Big Bend gambusia and its habitat according to State and Federal laws so they may properly identify the species and be aware of the potential threats and hazards to its continued existence.

# 5.2 Habitat integrity.

Agencies with jurisdiction over project activities which could modify the existing habitat in any way should be kept informed of the status of the Big Bend gambusia, its distribution and needs. Section 7 consultation requirements mandate that Federal project specifications preclude any adverse effect on listed species. Protection of the species is a joint responsibility of the U.S. Fish and Wildlife Service, the National Park Service, the State of Texas, and to an extent, the Republic of Mexico.

#### LITERATURE CITED

- Alvarez del Villar, Jose. 1970. Pecea Mexicanos (claves). Serie investigación pesquera, estudia No. 1. 166 pp.
- Brune, Gunnar. 1981. Springs of Texas. Vol. I. Branch-Smith, Inc. Fort Worth.
- Hubbs, Carl L. 1929. Studies of the fishes of the or&r Cyprinodontes VIII. Gambusia gaigei, a new species from the Rio Grande. occ. Pap. Mus. Zool. Univ. Mich. 198:1-11.
- Hubbs, Clark. 1959. High incidence of vertebral deformities in two natural populations of fishes inhabiting warm springs. Ecology 40:154-155.
- and Harold J. Broderick. 1963. Current abundance of <u>Gambusia gaigei</u>, an endangered fish species. Southwestern Naturalist 8:46-48.
- and Anthony A. Echelle. 1972. Endangered non-game fishes of the upper Rio Grande Basin. <u>In</u>: Endangered vertebrates in the Southwest. William C. Huey (ed.). New Mexico Department of Game and Fish: 147-167.
- Plan for Big Bend Gambusia, Big Bead National Park, Texas. National Park Service.
- Robert R. Miller, Robert J. Edwards, Kenneth W. Thompson, Edie Marsh,
  Gary P. Garrett, Gary L. Powell, D. J. Morris, and Robert W. Zerr. 1977b.
  Fishes inhabiting the Rio Grande, Texas-Mexico, between El Paso and the
  Pecos confluence. In: Importance, Preservation and Management of
  Riparian Habitat. A symposium. USDA Forest Service, General Technical
  Report. RM-43. R. Roy Johnson and Dale Jones (eds.). 91-97.
- , and Victor G. Springer. 1957. A revision of the Gambusia nobilis species group, with descriptions of three new species, and notes on their variation, ecology, and evolution. Texas J. Sci. 9:279-327.
- and John G. Williams. 1979. A review of circumstances affecting the abundance of <u>Gambusia gaigei</u>, an endangered fish endemic to Big Bend National Park. In: Proceedings **Firs** t Conference on Scientific Research in the National **Farks**, Vol. 1. Robert Linn (ed.). 631-635.
- Meffe, Gary K. 1983. Ecology of species replacement in the Sonoran topminnow (Poeciliopsis occidentalis) by mosquito fish (Gambusia affinis) in Arizona. Ph.d. dissertation. Acizona State Univ. Tempe, AZ

- Miller, Robert R. i978. Composition and derivation of the native fish fauna of the Chihuahusn desert region. In: Transactions of the Symposium on the Biological Resources of the Chihuahuan Desert Region, U.S.-Mexico. Roland H. Wauer and David H. Riskind (eds.), National Park Service, GPO, Washington, D.C.:365-381.
- Minckley, Wendell L. 1962. Two new species of fishes of the genus Gambusia (Poeciliidae) from northeastern Mexico. Copeia 1962: 391-396.
- Moore, G. 1980. Aquifer test of geothermal Spring 4, Rio Grande Village, Big Bend National Park, National Park Service, Water Resources Report No. 80-8. Southwest Regional Office, Santa Fe, NM 87501.
- Peden, A. E. 1973. Variation in anal spot expression of gambueine females and its effect on male courtship. Copeia 1973: 250-263.
- Rivas, L. R. 1963. Subgenera and species groups in the poeciliid fish genus Gambusia Poey. Copeia 1963: 331-347.
- Rosen, D. E. and R. M. Bailey. 1966. The poeciliid fishes (Cyprinodontiformes), their structure, zoogeography, and systematics. Bull. Amer. Mus. Nat. Hist. 126: 1-176.
- Wauer, Roland H. 1373. Naturalists Big Bend. Peregrine Prod., Santa Fe, New Mexico.

# PART III

# IMPLEMENTATION SCHEDULE

The table that follows is a summary of scheduled actions and costs for the Big Ben Gambusia Recovery Plan. This table indicates the priority in scheduling tasks to meet the objectives, which agencies are responsible for these tasks, a timetable for accomplishing them, and their estimated costs. Implementating Part III is the action of the recovery plan that, when accomplished, will satisfy the prime objective.

### GENERAL CATEGORIES FOR IMPLEMENTATION SCHEDULES

# Information Gathering I or R (Research) Acquis

- 1. Population status
- 2. Habitat statue
- 3. Habitat requirements
- 4. Management te chniques
- 5. Taxonomic studies
- 6. Demographic studies
- 7. Propagation
- 8. Migration
- 9. Predation
- 10. Competition
- 11. Di sease
- 12. Environmental contaminant
- 13. Reintroduction
- 14. Other information

# Acquisition -A

- 1. Lease
- 2. Easement
- 3. Management agreement
- 4. Exchange
- 5. Withdrawal
- 6. Fee title
- 7. Other

# Management - M

- 1. Propagation
- 2. Reintroduction
- 3. Habitat maintenance and manipulation
- 4. Predator and competitor control
- 5. Depredation contra 1
- 6. Disease control
- 7. Cther management

### Other -0

- 1. Inform& Ion and education
- 2. Law enforcement
- 3. Regulations
- 4. Administration

# Task Priority

Priority' 1 - An action that must be taken to prevent extinction or to prevent the species from declining irreversibly.

Priority 2 - An action that must be taken to prevent a significant decline in species population habitat quality or **some other significant** negative impact short of extinction.

Priority 3 - All other actions necessary to provide for full recovery of the species.

PART III - IMPLEMENTATION SCHEDULE

-	PLAN TASK		PRIORITY #	<u> </u>	RESPONSIBLE AGENCY			FISCAL YEAR COSTS *			COMMENTS
GENERAL		TASK #		TASK	FWS		OTHER		EST.)		
CATEGORY	(2)	(3)	(4)	DURATION (5)	EGION (6)	PROGRAM (6a)	(7)	N85 <b>(8)</b>	N86	N87	(9)
(1)	(2)	(3)	(4)	(3)	(0)	(Ua)	(/)	(8)			())
<b>R10 R11</b> <b>R9</b> R6 R3	Identify population needs	1.1	1	3 years	2	FR RES	NPS	1. <b>0</b> ,000	to ,000	LO, <b>000</b>	Consist of eubtaske 1.11 through 1.17
11 and <b>12</b>	<pre>Monitor populations and habitat</pre>	1.1	2	ongoing	2	SE	NPS	1. ,000	L ,000	1,000	
м3	Supplement spring flow	1.31	2	1 year	2	SE	NPS	2.5,000			
м3	Manage springs 1 and 4	1.32 1.	l <sub>3</sub> 2	ongoing	2	SE	NPS				
М3	Provide water to the campground from a well	1.34	3	<b>l</b> year	2		NPS				
M4	Eradicatre G. affinis from campground area	1.35	3	<b>l</b> year	2	FR SE	nps Tp&W	i!,000			
М3	Minimize campground impact	1.36	' 3	2 years	2	SE	NPS				
M2	Determine suitable stocking locations	2.1	3	1 year	2	SE	NPS TP&W	i! ,000			
M2	Stock Big Bend <b>Gabmusia</b>	2.2	3	ongoing as necessary	2	SE	nps <b>TP&amp;W</b>				
М7	Control mosquitos with Big Bend Gambusia	2.3	3	ongoing	2	SE	NPS				

<sup>\*</sup>Costs refer to USFWS expenditures only.

PART III - IMPLEMENTATION SCHEDULE

GENERAL CATEGORY _ (1) _	PLAN TASK (2)	'ASK #	'RIORITY #	TASK DURATION _ (5) _	RESPONSIBLE AGENCY			FISCAL YEAR <b>OSTS</b>			COMMENTS
					FWS REGION PROGRAM		OTHER	(EST.)* <b>FY85 FY86 F</b>		FY87	
					(6)	(6a)	(7)		1100		(9)
M2	Monitor relocated population8	1.4	3	ongoing	2	SE	NPS		assoc <b>3k 1.2</b>		
Ml	Maintain a captive population	3.0	2	ongoing	2	SE FR		<b>;</b> ,000	5,000	<b>;</b> ,000	
01	Prepare a information pamphlet	1	3	l year	2	SE	NPS	,000			
01	Conduct interpretive program	1.2	، 3	ongoing	. 2	SE	NPS				
02	Enforce all laws to protect the <b>species</b>	j	3	ongoing	2	LE	NPS	i ,000	5,000	<b>;</b> ,000	

<sup>\*</sup> Cost r fer to USPWS expenditures only

# PART IV - APPENDIX

#### APPENDIX I

## Replies to Comments

- A-l Recovery plan changed to put water needs of Big Bend gambusia ahead of campground domestic water use.
- A-2 Agreed and Recovery Plan changed accordingly.
- A-3 The Preface section of the recovery plan was transferred to the Disclaimer section and NPS added.
- A-4 Corrected.
- A-5 The paragraph was not changed since it adequately describes the taxonomic status of the species.
- A-6 Rewr it ten.
- A-7 Corrected.
- A-8 Sentence was rewritten.
- A-9 Corrected.
- A-10 Changed.
- A-11 Changed as suggested.
- A-12 Changed as suggested.
- A-13 A short discussion of threats has been Included.
- A-14 The Literature Cited section was changed to Miller (1978).
- A-15 The Recovery Team discussed the wording of the **Taxonomic** section of the recovery plan and decided that the section should remain as written because it **is** more accurate. Some changes for clarity have been added.
- A-16 Changed as suggested.
- A-17 Map changed as suggested.

- A-18 Discussion of future management goals added.
- A-19 The discussion of **the** primary objective was rewritten to expand upon recovery goals and habitat **limitations**.
- A-20 Added as suggested.
- A-21 Some generic information has been gained about <u>G. affinis</u> competition with <u>Gilatopulnnows</u> (<u>Paeciliopsis</u> <u>occidentalis</u>). That data has already been incorporated in this **plan**, but a great deal of **specific** knowledge is still lacking.
- A-22 Discussion added concerning potential for exotic fish introduction if pumping from the river occurs.
- A-23 Number is unknown at present. Investigation of potentials in various springs will have to be accomplished.
- A-2 4 Changed.
- A-25 Changed accordingly.
- A-26 The Implementation Schedule has been rewritten as suggested.
- **A-27** Changed as suggested.



### TEXAS PARKS AND WILDLIFE DEPARTMENT 4200 Smith School Road Austin, Tozas 78744

#### COMMISSIONERS

EDWIN L COX JR Chairman, Athens

GEORGE R. BOUN Vice-Chairman, nouston

WM. O. BRAECKLEIN

May 18, 1984

Mr. Michael J. Spear

Albuquerque, New Mexico

U. S. Fish and Wildlife Service

Director, Region 2

P. 0. Box 1306

Dallas

WM. I GRAHAM Amarillo

RICHARD R. MORRISON, III

Clear Lake city

W 8. OSBORN, JR Santa Elena

Dear Mike:

PERKINS O. SAMS Midland

DR RAY E. SANTOS Lubbock

WM M WHELESS, NI Houston

A - 1

A-2

This is in response to your letter of March 29, 1984

regarding the agency review draft recovery plan for the Big Bend gambusia.

87103

Overall, the recovery plan embodies strategies which, when implemented, should ensure survival of the species. The plan should succeed although, in view of the restricted distribution and problems for survival continually facing this species, recovery to the point of delisting or downlisting does not appear likely.

Our comments have been incorporated into the returned

Many of them relate to minor typographical errors, but two warrant additional discussion here. Recovery Outline Narrative 1.33 places the needs of campground domestic water above the needs of the Big Bend gambusia. It would seem appropriate in a fish species recovery plan that the needs of people would be subordinate. Another source of water for people should be sought--perhaps purified Rio Grande water. In 2.32, reference is made to recovery team approved Shouldn't this be <u>recommended</u> actions? actions.

Thank you for the opportunity to comment on this matter.

Sincerely,

Travis rles DJ Executive Director

CDT:FEP:aeh

Enclosure

**RECEIVED** BOF & W-REG. 2

> MAY22 1983 OFFICE OF THE REGIONAL DISCE SA

FWS REG 2 RECEIVED

JOHNSON

Burton Cortey

Halverson

PHES D. TRAVI

Energy Directo

Holin, "

**B**otanist

Норр

Padilla SANCHEZ

FILE

AHR.

PAO

EFO.

FILE

CL5./8

LE\_

LANGOWSKI



# United States Department of the Interior

Zulon III

# NATIONAL PARK SERVICE

Big Bend National Park
Big Bend National Park, Texas 79834

IN REPLY REFER TO:

N1621

May 14, 1384

Memorandum

Jo:

Assistant Regional Director, U.S.Fish and Wildlife Service,

Region 2

From:

Superintendent, Big Bend National Park

Subject: Agency Review Draft Recovery Plan for Big Bend Gambusia,

(Gambusia gaigei Hubbs 1929)

Qur comments on the Draft Recovery Plan are as follows:

A-3 1. Preface, paragraph 3, last sentence - Omit Texas Parks and Wildlife Department and insert National Park Service.

**A-4** 2. Page 1, Part I, Introduction, second paragraph, fourth line - Omit see.

 $A-5\ 3$ . Page 1, Part I, Taxonomic Status - This paragraph needs clarification.

4. Page 2, Part I, Distribution and Description of Habitat, first paragraph, first sentence - This sentence should read: The Big Bend Gambusia is known only from spring habitats in the vicinity of Equillas crossing and Rio Grande Village in Big Bend National Park.

**A-7** 5. Figure 2 - The Barkley Cabin is misspelled. Berkeley is the correct spelling.

A-8 6. Page 6, # 2, Development, second sentence - Omit one "in the future."

7. Page 10, paragraph 2, second line - Barkley House is misspelied. Berke'iey is the correct spelling.

We appreciate the opportunity to comment on the Draft Recovery Plan.

H. Gilbert Lusk



# United States Department of the Interior

FISH AND WILDLIFE SERVICE WASHINGTON, D.C. 20240

ADDRESS ONLY THE DIRECTOR, FISH AND WILDLIFE SERVICE

In Reply Refer Jo:

Memorandum

Jo:

Regional Director, Region 2 (ARD/AFF)

From:

Di rector

Subject:

Big Bend Gambusia Recovery Plan - Agency Review Draft

Attached is a copy of the subject plan with numerous editorial and The majority of these items substantive comments indicated in the margins. were brought to your attention in the Technical Review Draft. Please pay close attention to the comments noted on pages 2, 3, 6, 11, 13, 15, 17, 18, and 21 of the attached plan. Some of the more substantive comments are reiterated below:

A-10 1. Title page: The The title page should include only the title, publisher, Director's signature line, and the date

A-11<sup>2</sup>. Refer to the Recovery Planning Guidelines for the Disclaimer page: format of the Disclaimer page. A Disclaimer page should immediately follow the title page. The disclaimer paragraph used in the Preface (page i, paragraph 4) should begin the Disclaimer page. The proper acknowledgement for the plan should be indicated as follows:

> The Big Bend Gambusia Recovery Plan, dated \_\_ prepared by the U.S. Fish and Wildlife Service in cooperation with the Rio Grande Fishes Recovery Team.

The list of team members and consultants should be included on the Also, the Disclaimer page should identify the source Disclaimer page. of additional copies:

> Fish and Wildlife Reference Service 1776 E. Jefferson Street, 4th Floor Rockville, Maryland 20852 301/468-1737, ext. 231, or Toll Free 1-800-582-3421

FWS REG 2 RECEIVED

DKD\_

AA.

ANY

AHR. LE\_

PAO. .EEO.

2-FILE - C L -

End. Sp. R-2

NOSUHO! MIGOWS!! 96wman Burten Carley

**Fielvorson** Helfman

Cleell Stellerud

**Ectonist** Fore

7 SASICHEZ

JUN 8'84

SE

A-123. The Table of Contents should follow the Disclaimer page. To be consistent with other plans, the headings for the parts of the plan should be changed to: Part I - Introduction; Part II - Recovery; Part III - Implementation Schedule; Part IV - Appendix.

FWS. Region 2

JUN 7

AFF.

The Literature Cited section should be the last section of Part II and immediately precede Part III • Implementation Schedule. The Objective should be identifed in the Table of Contents.

- A-13 4. Page 1, Introduction: The introduction should include a brief discussion of the threats to the <u>Gambusia gaiqei</u>.
- A-14 5. Page 1, Taxonomic Status: The reference to Miller (1978) is listed as Miller (1977) in the Literature Cited section.
- **A-15** 6. Page 1, Taxonomic Status: The first two sentences are confusing. We suggest the following rewrite:

The Big Bend gambusia was described by **Hubbs** (1929). Although there has been some disagreement (Rosen and **Bailey**, 1966) regarding the synonomy of the species, the toncensus is that <u>Gambusia gaigei</u> is a valid species (Rivas, 1966; Peden, 1975; Aivarez, 1970; Miller, 1977).

- A-16 7. Page I, Taxonomic Status; This paragraph should be divided into two paragraphs: one on the synonomy discussion and another on the relationship, of <u>G. gaiqei</u> to the <u>G. senilis</u> species group.
- **A-17** 8. Figure I: The map would convey more information if the locations of present and extinct populations were labeled.
- A-18 9. Page 12: To complete the management scenario, briefly discuss future management galls.
- A-1910. Page 15, Part II: The discussion of the primary objective should include quantifiable goals with regard to the habitat and the species. Logically, the goals should correspond to the factors for which the Big Bend gambusia was listed.

If it appears likely the species cannot be reclassified or delisted, this should be clearly stated and the quantified goal for maintaining or preventing extinction of the species should be stated.

- **A-20** 11. Page 15, **Stepdown** Outline: Add the following **subtasks** under Task 2.0:
  - 2.1 Determine suitable locations
  - 2. 5 Manage introduced populations
- **A-21** 12. Page 17, Task 1.11: Since competition with G. affinis is a problem for most small listed fishes in Region 2, could this threat be studied on a more general basis and the data applied to more than one species?

- A-22 13. Page 20, Task 1.34: Please add some discussion on the necessity of this task. What will be enhanced by the accomplishment of this task?
- A-23 14. Page 21, Task 2.0: Approximately how many potential sites exist for reintroduction?
- A-24 15. Page 24, Appendix: Refer to the Recovery Planning Guidelines. The Appendix should be moved to the end of the plan, with a title page immediately preceding it.
- A-25 16. Implementation Schedule: Refer to the Recovery Planning Guidelines.
  Include a title page for the Implementation Schedule. The title page should include the definitions for the task priorities, general implementation categories, and abbreviations. Attached are examples from another plan.
- A-26 17. Implementation Schedule: The Implementation Schedule is not acceptable Recovery tasks should be identified as specifically as possible because this schedule will become the key for all Service activities (including funding recovery actions) involved in the recovery of the species. As you know, the review of permit proposals, Section 7 consultations, unsolicited proposals, State Federal Aid proposals, and all other funding requests will be examined against the recovery plan and corresponding Implementation Schedule. Subtasks must be included if the Implementation Schedule is to be useful.
- A-27 18. Implementation Schedule: Task 1.0 should be priority 1.

We hope these comments will be helpful in preparing the final draft. If you feel that any of these comments do not warrant revision of this draft, please provide your rationale, via return memorandum, prior to the Regional Director's approval. Upon approval, please send a copy of the signature page. Also, please send 30 copies of the printed plan when it is available.

Winds Deficiel

Attachments

#### APPENDIX II

# Monitoring Procedures

Sampling should be done biannually (February and August) with 3mm mesh seines; 3-meters long (pond) and 1.5-meters long (stream). Five seine hauls should be made at roughly equidistant points around the perimeter of the refugium pond. Each haul should begin 3-5 meters offshore and extend to the shore. Two seine hauls should be made in the pond outlet stream, one haul near the pond outlet, the other near the terminus. These hauls should be 2-3 meters long and should be made in the direction of s treamflow. The pond and ditch should be visually examined to ascertain presence of any other species. All seines used should be sun dried before use to preclude introduction 'of other fishes. Samples of 15 Big Bend gambusia (preferably males) should be preserved from both localities (pond and ditch) in separate labeled jars with 10% formalin. The preserved sample should be obtained from the first seine haul(s). If the fish are scarce, the number preserved may be reduced to avoid depletion of the population. All other fish captured should be visually examined to ensure that no other species are present; of particular concern is the mosquitofish, Gambusia affinis. Because of the difficulty of field separation of Gambusia species, selection of onsite supervisory personnel should be reviewed by the recovery team. The preserved specimens should be stored in an appropriate museum. The park should maintain a written log regarding time when collections were made, personnel involved, and disposition of samples. State and park collecting permits will be acquired before collections are made.